

INFLUENCE OF SUCROSE CONCENTRATION ON *IN VITRO* MULTIPLICATION OF *RIBES RUBRUM* SPECIES

Carmen Manole², Viorica Bălan², Cătălina Tudora¹, M. Buțu¹, Georgeta Fidler¹, Steliana Rodino¹, Diana Golea¹, Alina Buțu¹

¹National Institute of Research and Development for Biological Sciences, Department of Biotechnology, 0630031, Splaiul Independentei 296, Bucharest, Romania, e-mail: alina_butu@yahoo.com

²University of Agronomic Sciences and Veterinary Medicine, Faculty of Agriculture, Bucharest, Romania

Abstract. Sucrose is one of the major sources of carbon and energy to the culture media used for in vitro cultures of plants. In this article, was studied the influence of different sucrose concentrations on in vitro multiplication of the species *Ribes rubrum* (15 g/l, 30 g/l, 45 g/l, 60 g/l) on MS culture medium containing half macroelements supplemented with growth regulators BA (0.2 mg/l), IBA (0.01 mg/l) at pH=5.7–5.8. The highest multiplication rate was observed using the culture medium with a sucrose concentration of 45 g/l.

Key words: *Ribes rubrum*, sucrose concentration, micropropagation

Introduction

Ribes rubrum (red currant) is a rare species from the spontaneous flora, with importance in both food and medical industry.

It is valuable for its medicinal properties and commercial use in pharmaceutical and food industry and for the potential represented by leafs, fruits, seeds, buds and flowers.

The presence of sugars in the culture is an important factor of *in vitro* multiplication of plants [RICARDO *et al.*, 2004].

In plant tissue culture is needed a medium containing a permanent flow of carbohydrates to support their in vitro survival [BUTNARIU *et al.*, 2011]; the photosynthetic activity of tissues can be reduced by using a low light intensity, humidity and relatively high gas exchange [REZA *et al.*, 2009].

In vitro multiplication of plant material depends on many factors including the concentration and type of exogenous carbon source added in the environment, serving as an energy source and necessary for maintaining osmotic potential [TOUQUEER *et al.*, 2007].

One of these components is the concentration of sucrose which was found to be an important factor in the osmotic and photosynthetic potential and rooting response of plants [NASSER *et al.*, 2005].

The present study was conducted to determine the influence of sucrose concentration in culture medium on *in vitro* multiplication of the explants taken from red

currant.

Materials and method

MS culture medium [MURASHIGE *et al.*, 1962], half macronutrients content, was supplemented with BA (0.2 mg/l) and IBA (0.01 mg/l) at pH 5.7–5.8.

Concentrations of sucrose added to MS medium were MS I (15 g/l), MS II (30 g/l), MS III (45 g/l), MS IV (60 g/l).

We used flasks (capacity 50 ml) containing 15–20 ml of culture medium autoclaved, and then maintained for 30 minutes to UV light.

The study conducted on total 25 samples for each variant of medium inoculating one explant; the tubes were closed and kept in climate chamber at temperature between 23–25°C and a photoperiod of 16 hours (fluorescent light).

Explants consisted of shoot apex and nodal segments taken from young shoots of *Ribes rubrum* with a size of 0.3–0.5cm.

After three weeks of growth in climatic chamber, measurements were made compared with plantlets obtained in other medium variants.

Results

In this experiment, significant differences were noticed on plant height (Table 1).

Culture medium containing different sucrose concentrations present different values of plant height after three weeks from



starting the experiment.

The results of *in vitro* multiplication on MS I medium culture, showed that the nodal segments were not developed.

Table 1

Mean values for plant height (PH), assessed at different sucrose concentrations (15, 30, 45, 60 g/l) and number of leaves (NL), after three weeks from cultivation

Treatments	Means	
	PH(cm)	NL
MS I (15 g/l)	–	–
MS II (30 g/l)	1.0	2 leaves (0.2, 0.6 cm)
	0.33	2 primary leaves
	0.50	0.2 cm
	0.63	0.3 cm
	0.40	0.2 cm
MS III (45 g/l)	0.80	2 leaves (primary leaf, 0.3cm)
	0.52	2 leaves (0.3, 0.2 cm)
	0.71	2 leaves (0.2 cm)
	1.20	1 leaf (0.8 cm)
	0.82	1 leaf (0.7 cm)
MS IV (60 g/l)	0.60	3 leaves (0.2 cm)
	1.00	2 leaves (0.2 cm)
	1.02	2 leaves (0.4, 0.2 cm)
	0.50	2 primary leaves
	0.60	1 leaf (0.2 cm)

In other cases, it was showed the importance of increasing sucrose concentration on the multiplication of *Ribes* species (*Figure 1*).

During the experiment, we decided that the reference medium should be the one developed by Murashige and Skoog [MURASHIGE *et.al.* 1962], with optimum concentration of sucrose 30 g/l.

In *Figure 1*, we can see that in all cases except the first one, the increase in height of plantlets was directly proportional to increasing sucrose concentration.

In other studies, the results obtained by increasing the concentration of sucrose were found to be at the expense of those obtained using mannitol [GARCIA *et.al.* 2002].

After 21 days, plant height was between 0.33–1.20 cm (*Table 1*); lower limit is common in MS II (30 g/l). The greatest height of plantlets was recorded in the culture medium using MS III 45 g/l, PH=120 cm.

In terms of leaf growth, MS III and MS IV had a higher growth rate than using MS II. In addition, considering their size, the largest was recorded in the multiplication on MS medium III, the maximum diameter was 10 cm. regarding their number, more leaves per plantlet occurs using MS IV.




Figure 1. Plant height mean values (PH), according to sucrose concentration in the culture medium

Conclusions

Sucrose concentration influenced

growth and accumulation of biomass (fresh weight) of *Ribes* plantlets propagated *in vitro*.



The presence of 45 g/l sucrose in the culture medium was the most efficient treatment for increasing height and fresh weight of plants cultivated *in vitro*.

Acknowledgement

This work was supported by the CORE Program BIODIV no 09–360104/2009.

References

1. Butnariu, Monica *et al.* (2011). Antimicrobial and anti-inflammatory activities of the volatile oil compounds from *Tropaeolum majus L.* (Nasturtium). African Journal of Biotechnology 10 (31): 5900–5909.
2. Ricardo T. De Faria, Fabiana N. Rodrigues, Luciana Do V.R. Oliveira, Claudio Muller, **2004**–*In vitro Dendrobium nobile plant growth and rooting in different sucrose concentrations*, Horticultura Brasileira, v. 22, no.4, p.780–783.
3. Reza Bahmani, Omid Karami and Mansour Gholami, **2009**–*Influence of Carbon Sources and Their Concentrations on Rooting and Hyperhydricity of Apple Rootstock CM.106*, World Applied Sciences Journal, v.6 (11), p. 1513–1517.
4. Touqeer Ahmad, Nadeem Akhtar Abbasi, Ishfaq Ahmad Hafiz And Ansar Ali, **2007**–*Comparison of sucrose and sorbitol as main carbon energy sources in micropropagation of peach rootstock GF-677*, Pakistan Journal of Botany, v. 39 (4), p. 1269–1275.
5. Nasser S. Al-Khalifah, S. Hadi and F. Khan, **2005**–*Influence of Sucrose Concentration on in vitro Growth of Five Rose (Rosa hybrida L.) Cultivars*, Plant Tissue Cult., v. 15(1), p. 43–49;
6. Murashige T. and F. Skoog, **1962**–*A revised medium for rapid growth and bioassays with tobacco tissue cultures*, Physiol. Plant., v.15, p. 478–497.
7. J.L. Garcia, J. Troncoso, R. Sarmiento & A. Troncoso, **2002**–*Influence of carbon source and concentration on the in vitro development of olive zygotic embryos and explants raised from them*, Plant Cell, Tissue and Organ Culture, v. 69, p. 95–100.

Received: June 5, 2011
Accepted: August 27, 2011

